



## Statistical methods for the meta-analysis of full ROC curves

Oliver Kuß (joint work with Annika Hoyer)

Institut für Biometrie und Epidemiologie, Deutsches Diabetes-Zentrum (DDZ),  
Leibniz-Zentrum für Diabetes-Forschung an der Heinrich-Heine-Universität  
Düsseldorf

Mai 04, 2016, 16:00 s.t.

Department of Statistics, Ludwigstr. 33, Room 144, Munich

Meta-analyses and systematic reviews are the cornerstones of evidence based medicine and inform treatment, diagnosis, or prevention of individual patients as well as policy decisions in health care. Statistical methods for the meta-analysis of intervention studies are well established today. Meta-analysis for diagnostic accuracy trials, however, has been a vivid research area in recent years which is especially due to the increased complexity of diagnostic studies with their bivariate outcome of sensitivity and specificity. An even more increased complexity arises when single studies do not only report a single pair of sensitivity and specificity, but a full ROC curve with several pairs of sensitivity and specificity, each pair for a different threshold. Researchers frequently ignore this information and use only one pair of sensitivity and specificity from each study to arrive at meta-analytic estimates. Although methods to deal with the full information have been proposed [1-5], these are not without problems, e.g., they are two-step approaches where estimation uncertainty from the first step is ignored in the second step, the number of thresholds has to be identical across studies, or the concrete values of thresholds are ignored thus making impossible clinically relevant inference on sensitivity and specificity at given thresholds.

We propose two approaches for the meta-analysis of full ROC curves that use the information from all thresholds. The first approach simply expands the standard bivariate random effects model to a meta-regression model. The second approach uses the interpretation of an ROC curve as a bivariate time-to-event model for interval-censored data. This work is motivated by two systematic reviews on population-based screening for type 2 diabetes mellitus [6,7] which report on 38 single studies to assess the HbA1c as a diagnostic marker. Both reviews report only single pairs of sensitivity and specificity from each single study, but an intensified search yields 124 pairs of sensitivity and specificity for 26 different HbA1c thresholds from the 38 single studies.

## References

1. Kester AD, Buntinx F. Meta-analysis of ROC curves. *Med Decis Making*. 2000 Oct-Dec;20(4):430-9.
2. Dukic V, Gatsonis C. Meta-analysis of diagnostic test accuracy assessment studies with varying number of thresholds. *Biometrics*. 2003 Dec;59:936-46.
3. Hamza TH, Arends LR, van Houwelingen HC, Stijnen T. Multivariate random effects meta-analysis of diagnostic tests with multiple thresholds. *BMC Med Res Methodol*. 2009 Nov 10;9:73.
4. Martínez-Cambor P. Fully non-parametric receiver operating characteristic curve estimation for random-effects meta-analysis. *Stat Methods Med Res*. 2014 May 28.
5. Riley RD, Takwoingi Y, Trikalinos T, Guha A, Biswas A, Ensor J, Morris RK, Deeks JJ. Meta-Analysis of Test Accuracy Studies with Multiple and Missing Thresholds: A Multivariate-Normal Model. *Journal of Biometrics & Biostatistics* 2014 5:196.
6. Bennett CM, Guo M, Dharmage SC. HbA(1c) as a screening tool for detection of Type 2 diabetes: a systematic review. *Diabet Med*. 2007 Apr;24(4):333-43.
7. Kodama S, Horikawa C, Fujihara K, Hirasawa R, Yachi Y, Yoshizawa S, Tanaka S, Sone Y, Shimano H, Iida KT, Saito K, Sone H. Use of high-normal levels of haemoglobin A(1C) and fasting plasma glucose for diabetes screening and for rediction: a meta-analysis. *Diabetes Metab Res Rev*. 2013 Nov;29(8):680-92.